



# On new sequences converging towards the Euler–Mascheroni constant

Cristinel Mortici\*

Valahia University of Târgoviște, Department of Mathematics, Bd. Unirii 18, 130082 Târgoviște, Romania

## ARTICLE INFO

### Article history:

Received 6 August 2009

Received in revised form 11 January 2010

Accepted 11 January 2010

### Keywords:

Euler–Mascheroni constant

Speed of convergence

Approximations

## ABSTRACT

We propose new sequences containing a modified logarithmic term which converge to the Euler–Mascheroni constant faster than sequences known from the literature.

© 2010 Elsevier Ltd. All rights reserved.

## 1. Introduction

The Euler–Mascheroni constant  $\gamma = 0.57721566490153286 \dots$  is defined in mathematics as the limit of the sequence

$$\gamma_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln n$$

and it has numerous applications in many areas of pure and applied mathematics, such as analysis, theory of probability, special functions, applied statistics, or number theory. This constant is strangely ubiquitary and is probably the third most important constant in mathematics, after  $\pi$  and  $e$ . Nevertheless, the intimate nature of  $\gamma$  is very elusive; for example, it is not known if it is a rational number. A good part of its mystery comes from the fact that the known algorithms converging to  $\gamma$  are not very fast, at least, when they are compared to similar algorithms for  $\pi$  and  $e$ .

The sequence  $(\gamma_n)_{n \geq 1}$  converges very slowly toward its limit, like  $n^{-1}$  and in consequence, many authors are preoccupied to improve its speed of convergence. We have

$$\frac{1}{2(n+1)} < \gamma_n - \gamma < \frac{1}{2n} \quad (\text{Young})$$

(see, [1–3]) and other estimations can be found in [4–6].

Recently, DeTemple [7,8] had a clever idea to substantially improve the speed of convergence by introducing the sequence

$$R_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln \left( n + \frac{1}{2} \right).$$

It converges to  $\gamma$  like  $n^{-2}$ , since

$$\frac{1}{24(n+1)^2} < R_n - \gamma < \frac{1}{24n^2} \quad (\text{DeTemple}).$$

\* Tel.: +40 722727627; fax: +40 0245213382.

E-mail address: [cmortici@valahia.ro](mailto:cmortici@valahia.ro).

We define here a new class of sequences of the form

$$M_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln \frac{P(n)}{Q(n)},$$

where  $P, Q$  are polynomials having the leading coefficient equal to one and

$$\deg P - \deg Q = 1.$$

The classical sequence  $(\gamma_n)_{n \geq 1}$  is obtained in case  $P(n) = n, Q(n) = 1$ , while DeTemple's sequence  $(R_n)_{n \geq 1}$  is obtained for  $P(n) = n + \frac{1}{2}, Q(n) = 1$ .

Precisely, we introduce the sequences  $(v_n)_{n \geq 1}$  and  $(\mu_n)_{n \geq 1}$  by

$$v_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln \frac{n^2 + n + \frac{7}{24}}{n + \frac{1}{2}}$$

and

$$\mu_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln \frac{n^3 + \frac{3}{2}n^2 + \frac{227}{240}n + \frac{107}{480}}{n^2 + n + \frac{97}{240}}$$

which converge to  $\gamma$  like  $n^{-4}$ , respective  $n^{-6}$ , since

$$\lim_{n \rightarrow \infty} n^4 (v_n - \gamma) = -\frac{37}{5760} \quad \text{and} \quad \lim_{n \rightarrow \infty} n^6 (\mu_n - \gamma) = \frac{74\,381}{29\,030\,400}.$$

Our study is based on the following result, which gives a method for measuring the speed of convergence:

**Lemma 1.1.** *If  $(\omega_n)_{n \geq 1}$  is convergent to zero and there exists the limit*

$$\lim_{n \rightarrow \infty} n^k (\omega_n - \omega_{n+1}) = l \in \mathbb{R}, \quad (1.1)$$

with  $k > 1$ , then there exists the limit:

$$\lim_{n \rightarrow \infty} n^{k-1} \omega_n = \frac{l}{k-1}.$$

See [9] for the proof of this lemma and additional remarks. This result can be also used for constructing some asymptotic expansions, or to accelerate some convergences. See, e.g., [10–16,9].

We see from Lemma 1.1 that the speed of convergence of the sequence  $(\omega_n)_{n \geq 1}$  is as higher as the value  $k$  satisfying (1.1) is greater.

## 2. The results

First we define the sequence

$$v_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln \frac{n^2 + an + b}{n + c}$$

and we are interested to find the values  $a, b, c$  which provide the fastest sequence  $(v_n)_{n \geq 1}$ . First

$$v_n - v_{n+1} = -\frac{1}{n+1} - \ln \frac{n^2 + an + b}{n + c} + \ln \frac{(n+1)^2 + a(n+1) + b}{n+1+c},$$

and we are concentrated to compute a limit of the form (1.1). To do this, we consider the representation in power series:

$$\begin{aligned} v_n - v_{n+1} &= \left(c - a + \frac{1}{2}\right) \frac{1}{n^2} - \left(-a^2 - a + c^2 + c + 2b + \frac{2}{3}\right) \frac{1}{n^3} \\ &\quad - \left(a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4}\right) \frac{1}{n^4} \\ &\quad + \left(a^4 + 2a^3 - 4a^2b + 2a^2 - 6ab + a + 2b^2 - 4b - c^4 - 2c^3 - 2c^2 - c - \frac{4}{5}\right) \frac{1}{n^5} + O\left(\frac{1}{n^6}\right). \end{aligned} \quad (2.1)$$

According to Lemma 1.1, we have three parameters  $a, b, c$ , which produce the fastest convergence of the sequence from (2.1)

$$\begin{cases} c - a + \frac{1}{2} = 0 \\ -a^2 - a + c^2 + c + 2b + \frac{2}{3} = 0 \\ a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4} = 0, \end{cases}$$

namely, if  $a = 1$ ,  $b = \frac{7}{24}$ ,  $c = \frac{1}{2}$ .

Using [Lemma 1.1](#) and relation (2.1), we can state the following

**Theorem 2.1.** (i) If  $c - a + \frac{1}{2} \neq 0$ , then the speed of convergence of the sequence  $(v_n)_{n \geq 1}$  is  $n^{-1}$ , since

$$\lim_{n \rightarrow \infty} n^2 (v_n - v_{n+1}) = c - a + \frac{1}{2} \quad \text{and} \quad \lim_{n \rightarrow \infty} n (v_n - \gamma) = c - a + \frac{1}{2} \neq 0.$$

(ii) If  $c - a + \frac{1}{2} = 0$  and  $-a^2 - a + c^2 + c + 2b + \frac{2}{3} \neq 0$ , then the speed of convergence of the sequence  $(v_n)_{n \geq 1}$  is  $n^{-2}$ , since

$$\lim_{n \rightarrow \infty} n^3 (v_n - v_{n+1}) = -a^2 - a + c^2 + c + 2b + \frac{2}{3}$$

and consequently,

$$\lim_{n \rightarrow \infty} n^2 (v_n - \gamma) = \frac{1}{2} \left( -a^2 - a + c^2 + c + 2b + \frac{2}{3} \right) \neq 0.$$

(iii) If  $c - a + \frac{1}{2} = 0$ ,  $-a^2 - a + c^2 + c + 2b + \frac{2}{3} = 0$  and  $a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4} \neq 0$ , then the speed of convergence of the sequence  $(v_n)_{n \geq 1}$  is  $n^{-3}$ , since

$$\lim_{n \rightarrow \infty} n^4 (v_n - v_{n+1}) = a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4}$$

and consequently,

$$\lim_{n \rightarrow \infty} n^3 (v_n - \gamma) = \frac{1}{3} \left( a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4} \right) \neq 0.$$

(iv) If  $c - a + \frac{1}{2} = 0$ ,  $-a^2 - a + c^2 + c + 2b + \frac{2}{3} = 0$  and  $a - 3b - c - 3ab + \frac{3}{2}a^2 + a^3 - \frac{3}{2}c^2 - c^3 - \frac{3}{4} = 0$ , (equivalently with  $a = 1$ ,  $b = 7/24$ ,  $c = 1/2$ ), then the speed of convergence of the sequence  $(v_n)_{n \geq 1}$  is  $n^{-4}$ , since

$$\lim_{n \rightarrow \infty} n^5 (v_n - v_{n+1}) = -\frac{37}{1440} \quad \text{and} \quad \lim_{n \rightarrow \infty} n^4 (v_n - \gamma) = -\frac{37}{5760}.$$

For  $a = 1$ ,  $b = 7/24$ ,  $c = 1/2$ , the power series (2.1) becomes

$$v_n - v_{n+1} = -\frac{37}{1440n^5} + O\left(\frac{1}{n^6}\right),$$

which explains the assertion (iv) of [Theorem 2.1](#).

Now we define the sequence

$$\mu_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln \frac{n^3 + dn^2 + fn + g}{n^2 + hn + k}$$

and we are interested to find the values  $d, f, g, h, k$  which provide the fastest sequence  $(\mu_n)_{n \geq 1}$ . As above, we consider the representation in power series:

$$\begin{aligned} \mu_n - \mu_{n+1} &= \left( h - d + \frac{1}{2} \right) \frac{1}{n^2} + \left( d^2 + d - h^2 - h - 2f + 2k - \frac{2}{3} \right) \frac{1}{n^3} \\ &\quad - \left( d - 3f + 3g - h + 3k - 3df + 3hk + \frac{3}{2}d^2 + d^3 - \frac{3}{2}h^2 - h^3 - \frac{3}{4} \right) \frac{1}{n^4} \\ &\quad - \left( 4f - d - 6g + h - 4k + 4d^2f - 4h^2k + 6df - 4dg \right. \\ &\quad \left. - 6hk - 2d^2 - 2d^3 - d^4 - 2f^2 + 2h^2 + 2h^3 + h^4 + 2k^2 + \frac{4}{5} \right) \frac{1}{n^5} \end{aligned}$$

$$\begin{aligned}
& + \left( 5f - d - 10g + h - 5k - 5df^2 + 10d^2f + 5d^3f - 5d^2g + 5hk^2 \right. \\
& - 10h^2k - 5h^3k + 10df - 10dg + 5fg - 10hk - \frac{5}{2}d^2 - \frac{10}{3}d^3 - \frac{5}{2}d^4 \\
& \left. - d^5 - 5f^2 + \frac{5}{2}h^2 + \frac{10}{3}h^3 + \frac{5}{2}h^4 + h^5 + 5k^2 + \frac{5}{6} \right) \frac{1}{n^6} + o\left(\frac{1}{n^7}\right). \quad (2.2)
\end{aligned}$$

According to Lemma 1.1, we see that the fastest sequence  $(\mu_n)_{n \geq 1}$  is obtained in case when as many of the first coefficients of (2.2) is canceled. As we have now five parameters  $d, f, g, h, k$ , they produce the best result if and only if  $\alpha = \beta = \delta = \eta = \sigma = 0$ , where

$$\begin{aligned}
\alpha &= h - d + \frac{1}{2} \\
\beta &= d^2 + d - h^2 - h - 2f + 2k - \frac{2}{3} \\
\delta &= - \left( d - 3f + 3g - h + 3k - 3df + 3hk + \frac{3}{2}d^2 + d^3 - \frac{3}{2}h^2 - h^3 - \frac{3}{4} \right) \\
\eta &= - \left( 4f - d - 6g + h - 4k + 4d^2f - 4h^2k + 6df - 4dg \right. \\
&\quad \left. - 6hk - 2d^2 - 2d^3 - d^4 - 2f^2 + 2h^2 + 2h^3 + h^4 + 2k^2 + \frac{4}{5} \right) \\
\sigma &= \left( 5f - d - 10g + h - 5k - 5df^2 + 10d^2f + 5d^3f - 5d^2g + 5hk^2 \right. \\
&\quad - 10h^2k - 5h^3k + 10df - 10dg + 5fg - 10hk - \frac{5}{2}d^2 - \frac{10}{3}d^3 - \frac{5}{2}d^4 \\
&\quad \left. - d^5 - 5f^2 + \frac{5}{2}h^2 + \frac{10}{3}h^3 + \frac{5}{2}h^4 + h^5 + 5k^2 + \frac{5}{6} \right).
\end{aligned}$$

The solution is  $d = \frac{3}{2}, f = \frac{227}{240}, g = \frac{107}{480}, h = 1, k = \frac{97}{240}$ , which produces the best sequence  $(\mu_n)_{n \geq 1}$ . Using Lemma 1.1 and relation (2.2), we can state the following

**Theorem 2.2.** (i) If  $\alpha \neq 0$ , then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-1}$ , since

$$\lim_{n \rightarrow \infty} n^2 (\mu_n - \mu_{n+1}) = \alpha \quad \text{and} \quad \lim_{n \rightarrow \infty} n (\mu_n - \gamma) = \alpha \neq 0.$$

(ii) If  $\alpha = 0$  and  $\beta \neq 0$ , then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-2}$ , since

$$\lim_{n \rightarrow \infty} n^3 (\mu_n - \mu_{n+1}) = \beta \quad \text{and} \quad \lim_{n \rightarrow \infty} n^2 (\mu_n - \gamma) = \beta/2 \neq 0.$$

(iii) If  $\alpha = \beta = 0$  and  $\delta \neq 0$ , then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-3}$ , since

$$\lim_{n \rightarrow \infty} n^4 (\mu_n - \mu_{n+1}) = \delta \quad \text{and} \quad \lim_{n \rightarrow \infty} n^3 (\mu_n - \gamma) = \delta/3 \neq 0.$$

(iv) If  $\alpha = \beta = \delta = 0$  and  $\eta \neq 0$ , then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-4}$ , since

$$\lim_{n \rightarrow \infty} n^5 (\mu_n - \mu_{n+1}) = \eta \quad \text{and} \quad \lim_{n \rightarrow \infty} n^4 (\mu_n - \gamma) = \eta/4 \neq 0.$$

(v) If  $\alpha = \beta = \delta = \eta = 0$  and  $\sigma \neq 0$ , then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-5}$ , since

$$\lim_{n \rightarrow \infty} n^6 (\mu_n - \mu_{n+1}) = \sigma \quad \text{and} \quad \lim_{n \rightarrow \infty} n^5 (\mu_n - \gamma) = \sigma/5 \neq 0.$$

(vi) If  $\alpha = \beta = \delta = \eta = \sigma = 0$  (equivalently with  $d = \frac{3}{2}, f = \frac{227}{240}, g = \frac{107}{480}, h = 1, k = \frac{97}{240}$ ), then the speed of convergence of the sequence  $(\mu_n)_{n \geq 1}$  is  $n^{-6}$ , since

$$\lim_{n \rightarrow \infty} n^7 (\mu_n - \mu_{n+1}) = \frac{74\,381}{4\,838\,400} \quad \text{and} \quad \lim_{n \rightarrow \infty} n^6 (\mu_n - \gamma) = \frac{74\,381}{29\,030\,400}.$$

For  $d = \frac{3}{2}$ ,  $f = \frac{227}{240}$ ,  $g = \frac{107}{480}$ ,  $h = 1$ ,  $k = \frac{97}{240}$ , the power series (2.2) becomes

$$\mu_n - \mu_{n+1} = \frac{74381}{4838400n^7} + O\left(\frac{1}{n^8}\right),$$

which explains the assertion (vi) of Theorem 2.2.

### 3. Concluding remarks

As least theoretically, further sequences of the form

$$M_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n} - \ln \frac{P(n)}{Q(n)}$$

can be defined, with  $\deg P = k + 1$ ,  $\deg Q = k \geq 3$ . As above,

$$M_n - M_{n+1} = -\frac{1}{n+1} - \ln \frac{P(n)Q(n+1)}{Q(n)P(n+1)} \quad (3.1)$$

and if we develop (3.1) in power series of  $n^{-1}$ , then the  $2k+1$  coefficients of the polynomials  $P$  and  $Q$  are the unique solution of the system obtained by imposing that the first  $2k+1$  coefficients of (3.1) vanish. In this case,

$$M_n - M_{n+1} = \frac{\theta}{n^{2k+3}} + O\left(\frac{1}{n^{2k+4}}\right),$$

with  $\theta \neq 0$ . From Lemma 1.1,  $(M_n)_{n \geq 1}$  tends to zero as  $n^{-(2k+2)}$ , since

$$\lim_{n \rightarrow \infty} n^{2k+2} M_n = \frac{\theta}{2k+2} \neq 0.$$

Finally, we offer some numerical computations to prove the superiority of our sequences  $(v_n)_{n \geq 1}$  and  $(\mu_n)_{n \geq 1}$  over the classical sequence  $(\gamma_n)_{n \geq 1}$  and the DeTemple sequence  $(R_n)_{n \geq 1}$ . As an example, we see that already  $\mu_1$  approximates  $\gamma$  with eight exact decimals.

$n$	$\gamma_n - \gamma$	$R_n - \gamma$	$\gamma - v_n$	$\mu_n - \gamma$
10	$4.9167 \times 10^{-2}$	$3.7733 \times 10^{-4}$	$5.2565 \times 10^{-7}$	<b><math>1.8859 \times 10^{-9}</math></b>
50	$9.9667 \times 10^{-3}$	$1.6337 \times 10^{-5}$	$9.8744 \times 10^{-10}$	<b><math>1.5438 \times 10^{-13}</math></b>
100	$4.9917 \times 10^{-3}$	$4.1252 \times 10^{-6}$	$6.2964 \times 10^{-11}$	<b><math>2.4863 \times 10^{-15}</math></b>
300	$1.6657 \times 10^{-3}$	$4.6142 \times 10^{-7}$	$7.8777 \times 10^{-13}$	<b><math>3.4796 \times 10^{-18}</math></b>

### Acknowledgements

The author thanks the editor and anonymous referees for their useful comments and suggestions.

### References

- [1] C. Mortici, An ultimate extremely accurate formula for approximation of the factorial function, Arch. Math. (Basel) 93 (1) (2009) 37–45.
- [2] C. Mortici, A. Vernescu, Some new facts in discrete asymptotic analysis, Math. Balkanica 21 (3–4) (2007) 301–308.
- [3] R.M. Young, Euler's constant, Math. Gaz. 75 (1991) 187–190.
- [4] H. Alzer, Inequalities for the gamma and polygamma functions, Abh. Math. Sem. Univ. Hamburg 68 (1998) 363–372.
- [5] G.D. Anderson, R.W. Barnard, K.C. Richards, M.K. Vamanamurthy, M. Vuorinen, Inequalities for zero-balanced hypergeometric functions, Trans. Amer. Math. Soc. 347 (5) (1995) 1713–1723.
- [6] S.R. Tims, J.A. Tyrrell, Approximate evaluation of Euler's constant, Math. Gaz. 55 (391) (1971) 65–67.
- [7] D.W. DeTemple, A quicker convergence to Euler's constant, Amer. Math. Monthly 100 (5) (1993) 468–470.
- [8] D.W. DeTemple, A geometric look at sequences that converge to Euler's constant, College Math. J. 3 (2006) 128–131.
- [9] C. Mortici, New approximations of the gamma function in terms of the digamma function, Appl. Math. Lett. 23 (1) (2010) 97–100.
- [10] C. Mortici, Product approximations via asymptotic integration, Amer. Math. Monthly 117 (5) (2010) 434–441.
- [11] C. Mortici, A. Vernescu, An improvement of the convergence speed of the sequence  $(\gamma_n)_{n \geq 1}$  converging to Euler's constant, An. Ştiinţ. Univ. "Ovidius" Constanţa 13 (1) (2005) 97–100.
- [12] C. Mortici, Completely monotonic functions associated with gamma function and applications, Carpathian J. Math. 25 (2) (2009) 186–191.
- [13] C. Mortici, New sharp bounds for gamma and digamma functions, An. Ştiinţ. Univ. Al. I. Cuza Iaşi Mat. LVI (2) (2010) (in press).
- [14] C. Mortici, Improved convergence towards generalized Euler–Mascheroni constant, Appl. Math. Comput. 215 (9) (2010) 3443–3448.
- [15] C. Mortici, Optimizing the rate of convergence in some new classes of sequences convergent to Euler's constant, Anal. Appl. (Singap.) 8 (1) (2010) 99–107.
- [16] C. Mortici, The proof of Muqattash–Yahdi conjecture, Math. Comput. Modelling (2010) doi:10.1016/j.mcm.2009.12.030.